

WAS THE UNIVERSE REIONIZED BY MASSIVE METAL FREE STARS?

J. STUART B. WYITHE¹ AND ABRAHAM LOEB²
 swyithe@isis.ph.unimelb.edu.au; loeb@sns.ias.edu
Draft version February 2, 2008

ABSTRACT

The *WMAP* satellite has measured a large optical depth to electron scattering after cosmological recombination of $\tau_{\text{es}} = 0.17 \pm 0.04$, implying significant reionization of the primordial gas only ~ 200 million years after the big bang. However, the most recent overlap of intergalactic HII regions must have occurred at $z \lesssim 9$ based on the Ly α forest constraint on the thermal history of the intergalactic medium. Here we argue that a first generation of metal-free stars with a heavy (rather than Salpeter) mass function is therefore *required* to account for much of the inferred optical depth. This conclusion holds if feedback regulates star formation in early dwarf galaxies as observed in present-day dwarfs.

Subject headings: cosmology: theory – early universe – intergalactic medium – Stars: formation

1. INTRODUCTION

The first stars in the universe (Pop-III) formed out of metal-free gas, relic from the big bang. The bottom-up hierarchy of structure formation in Cold Dark Matter (CDM) models implies that stars started to form in CDM halos just above the cosmological Jeans mass of $\sim 10^5 M_\odot$ at redshifts as high as $z \sim 20 - 30$ (see review by Barkana & Loeb 2001 and references therein). Simulations of metal-free star formation indicate that the first stars had high masses ($M \gtrsim 100 M_\odot$) since gas cooling by molecular hydrogen (H_2) cannot lower the gas temperature below $\sim 200\text{K}$ (Bromm, Coppi, & Larson 2002; Abel, Bryan & Norman 2002). Massive, metal-free stars shine at their Eddington luminosity, $L_E \propto M$, and have roughly constant effective (surface) temperatures of $\sim 10^5\text{K}$ and lifetimes of $\sim 3 \times 10^6\text{yr}$, independent of their mass (Bromm, Kudritzki, & Loeb 2001). This implies that the number of ionizing photons produced per baryon incorporated into these stars ($\sim 4.5 \times 10^4$) was independent of their mass function, and larger by more than an order of magnitude than the yield of the observed Pop-II (low-mass, metal-rich) stars.

Did the first generation of metal-free stars produce a sufficient total number of ionizing photons to reionize the universe? This question is difficult to answer theoretically. The UV radiation from the massive stars can easily dissociate the fragile intergalactic H_2 molecules which are responsible for star formation in mini-halos with virial temperatures $T_{\text{vir}} \lesssim 10^{2.4}\text{K}$ (Haiman, Rees, & Loeb 1997; Haiman, Abel & Rees 2000; Ricotti et al. 2002). Moreover they could evaporate the gas out of the shallow potential wells of the first CDM mini-halos (Barkana & Loeb 1999; Nishi & Tashiro 2000). Finally the hydrodynamic energy released by a single supernova explosion is much larger than the gravitational binding energy of the gas in these halos (Barkana & Loeb 2001). If feedback quenches star formation in mini-halos, then Pop-III star formation will be dominated by more massive halos where H_2 can form in dense gas shielded from the UV background (Oh & Haiman 2002; Mackey, Bromm & Hernquist 2002).

Analysis of the first year data from the *WMAP* satel-

lite suggests a large optical depth to electron scattering $\tau_{\text{es}} = 0.17 \pm 0.04$, implying significant reionization at redshifts as high as $\sim 17 \pm 5$ (Kogut et al. 2003; Spergel et al. 2003). While this result is not surprising in view of earlier detailed calculations of the reionization history by metal-free stars (Wyithe & Loeb 2003; Paper I), it offers an intriguing *empirical* path for answering the aforementioned question. In this *Letter*, we argue that metal-free stars are *required* by the *WMAP* measurement of τ_{es} .

We assume the most recent cosmological parameters obtained through fits to *WMAP* data (Spergel et al. 2003). These include density parameters of $\Omega_m = 0.27$ in matter, $\Omega_b = 0.044$ in baryons, $\Omega_\Lambda = 0.73$ in a cosmological constant, a Hubble constant of $H_0 = 71 \text{ km s}^{-1} \text{ Mpc}^{-1}$, an rms amplitude of $\sigma_8 = 0.84$ for mass density fluctuations in a sphere of radius $8h^{-1}\text{Mpc}$, and a primordial power-spectrum with a power-law index $n = 1$. Any tilt in the power spectrum at the level that is marginally indicated by the combination of the *WMAP* data with other data sets (Spergel et al. 2003) would reduce the small-scale power, delay reionization, and strengthen our conclusion.

2. OPTICAL DEPTH TO ELECTRON SCATTERING

Different reionization histories result in different large-scale polarization anisotropies of the microwave background (Zaldarriaga 1997; Kaplinghat et al. 2002). A simple measure of the reionization history is the integrated optical depth to electron scattering $\tau_{\text{es}} = \int_0^{10^3} dz |cdt/dz| \sigma_T n_e(z)$, where σ_T is the Thomson cross-section and $(dt/dz)^{-1} = -H_0(1+z)\sqrt{\Omega_m(1+z)^3 + \Omega_\Lambda}$. The electron number density, n_e , depends on the mass filling factors Q^m of the different ionized species in the gas,

$$n_e = (1+z)^3 [Q_{\text{H}^+}^m n_{\text{H}}^0 + Q_{\text{He}^+}^m n_{\text{He}}^0 + 2Q_{\text{He}^{++}}^m n_{\text{He}}^0], \quad (1)$$

where n^0 denotes the total present-day number densities of hydrogen (H), and helium (He).

3. REIONIZATION HISTORIES

We now consider reionization histories under different model assumptions. The histories follow the reionization

¹ The University of Melbourne, Parkville, Vic, Australia

² Institute for Advanced Study, Princeton, NJ 08540; Guggenheim Fellow; On leave from the Astronomy Department, Harvard University

of hydrogen as well as the single and double reionization of helium in an inhomogeneous IGM (Miralda-Escude et al. 2000). We include helium in our calculations since the harder spectrum of massive Pop-III stars can reionize both hydrogen and helium at high redshift (Oh et al. 2001; Venkatesan, Tumlinson & Shull 2003).

We calculate the emissivity of quasars described by the luminosity function of Wyithe & Loeb (2002a) as well as the emissivity of high redshift zero-metallicity stars and Pop-II stars (the latter is discussed in § 3.1). Our model has two free parameters: (i) the transition redshift, z_{tran} , at which Pop-II stars start to dominate the cosmic radiation field of ionizing photons; and (ii) the product of the star formation efficiency in massive galaxies f_* (=stellar mass/total initial baryonic mass) and the escape fraction f_{esc} of ionizing photons from galaxies. In Paper I we presented contours of τ_{es} as a function of z_{tran} and $f_* f_{\text{esc}}$. We have found that if the star formation efficiency in early low-mass galaxies resembled that in local dwarf galaxies with the same circular velocity, then values of $f_* f_{\text{esc}} \sim 0.05$ – 0.2 and $z_{\text{tran}} \sim 15$ lead to τ_{es} within the range measured by *WMAP*. Below we summarize the basic ingredients of our model; more details can be found in Paper I.

We treat the IGM as a two-phase medium. In regions of reionized IGM, star formation can only be initiated in halos that collapse above a virial temperature threshold of $T_{\text{min}} = 2.5 \times 10^5 \text{K}$, corresponding to the cosmological Jeans mass of photo-ionized IGM at $\sim 10^4 \text{K}$ (e.g., Thoul & Weinberg 1996; Kitayama & Ikeuchi 2000). In the neutral IGM, star formation is limited to halos with virial temperatures above the minimum for efficient gas cooling. If only atomic hydrogen is present, then $T_{\text{min}} = 10^4 \text{K}$. However if molecular hydrogen is present then $T_{\text{min}} = 10^{2.4} \text{K}$ (Haiman, Thoul, & Loeb 1996; Haiman, Abel, & Rees 2000) and star formation can proceed in much smaller halos. We begin by discussing the case with $T_{\text{min}} = 10^4 \text{K}$.

3.1. Ionizing Photons from Stars

To compute the comoving densities of stellar ionizing photons (n_γ) we integrate over model spectra for stellar populations to find the total number of ionizing photons (N_γ) emitted per baryon incorporated into stars. We find

$$\frac{dn_\gamma}{dz} = N_\gamma f_{\text{esc}} \frac{dF_*}{dz} n_b, \quad (2)$$

where n_b is the number density of baryons, and $F_*(z)$ is the fraction of baryons in the universe at redshift z that collapsed, cooled, and formed stars.

A recent analysis of SDSS data (Kauffmann et al. 2002) has shown that in a large sample of local galaxies, the ratio between the total stellar (M_*) and dark matter halo (M) masses scales as $M^{2/3}$ for $M_* < 3 \times 10^{10} M_\odot$, but is constant for larger stellar masses. Thus the star formation efficiency within dwarf galaxies is proportional to $M^{2/3}$. Since star formation is thought to be regulated by supernova feedback (Dekel & Silk 1986; Dekel & Woo 2002), and the supernova energy output is proportional to M_* while the gravitational binding energy scales as $M v_{\text{cir}}^2 \propto M^{5/3}$, more massive galaxies retain a

larger fraction of their gas. Using the stellar mass Tully-Fisher relation of Bell & De Jong (2001), we find the threshold circular velocity $v_* = 176 \text{ km s}^{-1}$ that corresponds to a stellar mass of $3 \times 10^{10} M_\odot$ at $z = 0$. The mass of a dark matter halo with circular velocity v_* is $M_{\text{halo}}^* \sim 5 \times 10^{10} [(1+z)/10]^{-3/2} M_\odot$. We define f_* as the star formation efficiency in galaxies with circular velocities larger than v_* ; below v_* the star formation efficiency is $\eta_* = f_* (M/M_{\text{halo}}^*)^{2/3}$. Given T_{min} we find

$$\frac{dF_*(T_{\text{min}})}{dz} = \frac{d}{dz} \frac{1}{\rho_m} \int_{M(T_{\text{min}})}^\infty dM \epsilon f_* M \frac{dn_{\text{st}}}{dM}, \quad (3)$$

where $\epsilon = 1$ for $M > M_{\text{halo}}^*$ and $\epsilon = (M/M_{\text{halo}}^*)^{2/3}$ for $M < M_{\text{halo}}^*$, ρ_m is the mean comoving mass density of matter, and $n_{\text{st}}(> M)$ is the comoving number density of halos with masses $> M$ (Sheth & Thormen 1999). Finally, the value of dF_*/dz in the two phase IGM required for calculation of dn_γ/dz may then be computed from

$$\frac{dF_*}{dz} = Q_{\text{H}^+}^m \frac{dF_*(2.5 \times 10^5 \text{K})}{dz} + (1 - Q_{\text{H}^+}^m) \frac{dF_*(10^4 \text{K})}{dz}. \quad (4)$$

3.2. The Need for Metal-Free stars

We begin with calculations of reionization histories in the absence of metal-free stars at all redshifts. We have calculated reionization histories for several values of $f_* f_{\text{esc}}$ in terms of the volume filling factor Q of ionized regions in the low overdensity ($\lesssim 20$) IGM (Miralda-Escude et al. 2000; Paper I), and computed the resulting values of τ_{es} .

Recently, Theuns et al. (2002) have argued that if HI reionization ended before $z = 9$, and the universe remained reionized after that, then adiabatic cooling would have lead to an IGM temperature at $z = 3.5$, much lower than inferred from the Ly α forest. As a result, the most recent overlap of intergalactic HII regions must have occurred at $z \lesssim 9$, unless there was significant photo-heating at $z \gg 3.5$ due to the double reionization of helium. In the absence of metal free stars we find that this constraint requires $f_* f_{\text{esc}} \lesssim 0.1$, resulting in a value of $\tau_{\text{es}} \lesssim 0.09$ that is not consistent with the *WMAP* data. On the other hand, if we neglect the constraint that HII overlap occurred at $z \lesssim 9$, we find that in the absence of metal-free stars, reionization histories resulting in $\tau_{\text{es}} \gtrsim 0.13$ imply HII overlap redshifts of ~ 15 . However assuming feedback regulation of star formation in low mass galaxies, these histories require the unphysical value of $f_* f_{\text{esc}} = 1.0$ in galaxies with $v_{\text{cir}} > v_*$. We therefore conclude that reionization by an early population of massive metal-free stars (with very large³ N_γ) must have contributed significantly to τ_{es} at $z \sim 15$.

The formation of metal free stars is moderated with cosmic time by the gradual metal enrichment of the IGM. It has been argued that the dominant mode of star formation change to Pop-II once the metallicity of the associated gas rises above $10^{-4} Z_\odot$ (Schneider et al. 2002; Bromm et al. 2001). If all Pop-III stars formed in the mass range of $140 M_\odot - 260 M_\odot$ where pair-unstable supernovae result in expulsion of metals into the surrounding gas (Heger & Woosley 2001), and if all metals were mixed

³ We conservatively adopt the ionizing photon yield of $100 M_\odot$ metal-free stars. The yield is larger by a factor of ~ 2 for Pop-III stars with masses $\gtrsim 300 M_\odot$ (Bromm, Kudritzki, & Loeb 2001). The use of the higher yield would have lead to the same τ_{es} for values of $f_* f_{\text{esc}}$ at $z > z_{\text{tran}}$ that are smaller by a factor of 2 than we quote.

uniformly throughout the entire IGM, then Pop-III star formation would have ceased as soon as ~ 0.3 ionizing photons were produced per cosmic hydrogen atom (Ciardi, Ferrara & White 2003). However, the highly inhomogeneous nature of the enrichment process (Furlanetto & Loeb 2003) and the formation of Pop-III stars beyond the above mass range, naturally lead to reionization.

Must the Pop-III stars possess a heavy initial mass function (IMF)? Tumlinson & Shull (2000) computed the ionizing photon output per baryon of zero-metallicity stars with a Salpeter IMF and found only a 50% increase over the values for metal enriched Pop-II stars with the same IMF. This moderate increase is to be compared with the increase by more than an order of magnitude for the massive Pop-III stars described by Bromm, Kudritzki, & Loeb (2001) (the number of ionizing photons produced per baryon is nearly independent of stellar mass). The arguments presented above therefore imply that Pop-III stars were very massive rather than of a Salpeter IMF. This provides the first observational support to the results from hydrodynamic simulations of early star formation (Bromm et al. 2002; Abel et al. 2002).

In generating the star formation history we have extrapolated results from the local galaxy sample, assuming that feedback governs the observed stellar mass as a function of halo potential depth. However high redshift galaxies may have cooling times that are shorter than the dynamical time of their cold gas. This may lead to a starburst that might render the time-delayed supernovae feedback less effective. We now briefly consider this case. Assuming the star formation efficiency to be independent of halo mass (case A in Paper I), we find that reasonable values of $f_\star f_{\text{esc}} \sim 0.1$ can yield $\tau_{\text{es}} \sim 0.17$ in the absence of massive Pop-III stars. However, the corresponding reionization histories also have an early HII overlap. Therefore, unless the global product of escape fraction and star formation efficiency was more than an order of magnitude larger at $z \sim 15$ than at $z \sim 9$, our conclusion that an early generation of massive metal free stars is required to produce both $\tau_{\text{es}} \sim 0.17$ and HII overlap at $z \lesssim 9$ is independent of whether feedback regulates star formation in low mass galaxies or not.

3.3. Reionization Histories

We now show that the joint constraints of $\tau_{\text{es}} = 0.17 \pm 0.04$ and HII overlap at $z \lesssim 9$ can be met simultaneously for values of $z_{\text{tran}} > 9$ and $f_\star f_{\text{esc}} < 1$ if massive metal-free stars were present at high redshifts. In Figure 1 we show two cases of the evolution of Q for hydrogen (solid lines), singly ionized helium (dashed lines) and doubly ionized helium (dotted lines). We assume⁴ $z_{\text{tran}} = 12$ and $f_\star f_{\text{esc}} = 0.08$ at $z < z_{\text{tran}}$, yielding the most recent HII overlap at $z \sim 9$. At $z > z_{\text{tran}}$ we assume $f_\star f_{\text{esc}} = 0.08$ (left) and $f_\star f_{\text{esc}} = 0.33$ (right). The latter allows for the possibility of a higher star formation efficiency⁵ or escape fraction (Steidel et al. 2001) at high redshift. In the former case there is a significant reionization but *failed* HII overlap at early times, followed by a recombination and a successful HII overlap at $z \sim 9$ due to Pop-II stars. The

two examples yield τ_{es} within one standard deviation and at the most likely value of *WMAP*, respectively. Note that our model leads to the double reionization of helium by quasars at $z \sim 3.5$ independent of the contribution from Pop-III stars. This is in excellent agreement with current observational data [e.g. Theuns et al. (2002) and references therein]. The reionization of HeIII by quasars is characterized by a sharp rise in our model, with $Q_{\text{He}^{++}}$ increasing from $\sim 50\%$ at $z \sim 4.3$ to $\sim 100\%$ at $z \sim 3.7$. We do not expect pre-heating from the reionization of HeIII to impact on the conclusion that the most recent HII overlap took place at $z \lesssim 9$.

Figure 1 also shows the evolution of Q for hydrogen when helium is ignored (light solid line). In these calculations all photons blueward of 13.6eV reionize hydrogen. We find that the additional ionizing photons result in overlap of hydrogen at an earlier epoch, but the net optical depth to electron scattering is decreased.

So far we have assumed that atomic cooling dictates the mass of the smallest galaxies. However it has been suggested that significant production of X-rays could catalyze molecular hydrogen production allowing star formation in mini-halos before reionization (Haiman, Rees, & Loeb 1996; Cen 2002). The light dashed lines in the left hand panels of Figure 1 show the evolution in Q for hydrogen (in the absence of helium) assuming $T_{\text{min}} = 10^{2.4}\text{K}$ in the neutral IGM. We find that the feedback-regulation built into the model sufficiently limits the star formation efficiency in mini-halos so that they do not contribute significantly to the reionization history.

4. CONCLUSIONS

We have shown that Pop-II stars alone cannot reionize the universe sufficiently early to produce the value of $\tau_{\text{es}} \sim 17 \pm 4\%$ inferred from the *WMAP* data (Kogut et al. 2003; Spergel et al. 2003). This conclusion holds if feedback regulates star formation in early low-mass galaxies as observed in present-day dwarf galaxies. Without metal free stars, the condition that the most recent overlap of intergalactic HII regions must have occurred at $z \lesssim 9$ (Theuns et al. 2002) generically yields $\tau_{\text{es}} \lesssim 9\%$. The joint requirements that $\tau_{\text{es}} \sim 17\%$ and that HII regions overlapped at $z \lesssim 9$ imply that metal-free stars reionized a significant fraction of the IGM at $z \sim 15$. Moreover, we find that these stars must have been *massive* to provide the required order-of-magnitude increase in the production rate of ionizing photons relative to low mass Pop-II stars (Bromm, Kudritzki & Loeb 2001). Pop-III stars with a Salpeter IMF provide only a 50% increase in N_γ (Tumlinson & Shull 2000) and cannot account for the large τ_{es} .

For plausible reionization histories with metal-free stars we have found that singly and doubly ionized helium contributes as much as $\Delta\tau_{\text{es}} = 0.02$, and should be included in reionization calculations. We also found that if the star formation efficiency is regulated in dwarf galaxies as observed in the local universe, then there is no significant contribution to the cosmic ionizing radiation field from halos with virial temperatures $T_{\text{vir}} \lesssim 10^4\text{K}$, even if molecular hydrogen permits cooling of gas inside these mini-halos.

⁴ Salvaterra & Ferrara (2003) have recently presented the observational case that $z_{\text{tran}} \lesssim 9$ through fitting the near-infrared background. The lower value of z_{tran} would allow a second HII overlap near $z \sim 7$ to be compatible with the large value of τ_{es} and IGM temperature constraints.

⁵ Note that the star formation efficiency in the low-mass galaxies of interest, $\eta_\star = 0.02f_\star (M/10^8 M_\odot)^{2/3} [(1+z)/15]$, is generally $\ll f_\star$.




FIG. 1.— Sample reionization histories assuming feedback regulated star formation, that produce large values of τ_{es} . Dark lines represent the evolution of the volume filling factors of ionized regions $Q_{\text{H}+}$, $Q_{\text{He}+}$ and $Q_{\text{He}++}$. We also show calculations of $Q_{\text{H}+}$ in the absence of helium (light lines). The solid light lines show results for a pre-reionization minimum virial temperature for galaxies of $T_{\text{min}} = 10^4 \text{K}$ (atomic cooling) while the dashed light line corresponds to $T_{\text{min}} = 10^{2.4} \text{K}$ (molecular cooling). Each example is labeled by the values of $f_* f_{\text{esc}}$ at $z < z_{\text{tran}}$ (and in parenthesis at $z > z_{\text{tran}}$), z_{tran} and τ_{es} .

The evolution of the ionization state and metal enrichment of the IGM during the reionization epoch can be best probed by infrared spectroscopy of gamma-ray burst (GRB) afterglows (Furlanetto & Loeb 2002). These transient sources should be both abundant (Bromm & Loeb 2002) and detectable (Lamb & Reichart 2000; Ciardi & Loeb 2000) out to high redshifts. The *Swift* satellite, planned for launch this year, is expected to detect ~ 100 GRBs per year, out of which $\gtrsim 20\%$ may originate at $z \gtrsim 5$ (Bromm & Loeb 2002). Photometric identification of the Ly α trough can be used to select the high-redshift afterglows for which spectroscopic follow-up observations may be conducted on a 10-m class telescope (Loeb 2001).

After the submission of this paper, a number of different groups have also commented on the implications of the large τ_{es} measured by *WMAP*. Some studies find that Pop-II stars are able to reionize the universe sufficiently early to produce $\tau_{\text{es}} \gtrsim 0.13$ (Ciardi, Ferrara & White 2003; Somerville & Livio 2003); however, this conclusion is challenged by the existence of the observed Gunn-Peterson

trough in quasars at $z \sim 6.3$ (as noted by Ciardi et al. 2003) and by the temperature evolution of the IGM (Theuns et al. 2002; Hui & Haiman 2003). Other authors agree that the combined requirements of a large τ_{es} and a late HII overlap implies that the efficiency of production of ionizing photons was higher at redshifts $z \gtrsim 10$ (Haiman & Holder 2003; Sokasian, Abel, Hernquist & Springel 2003; Cen 2003). The conclusion that Pop-III stars needed to be massive to produce the observed τ_{es} was also reached by Cen (2003) in agreement with our results.

We thank Rennan Barkana, Volker Bromm, Steve Furlanetto, Rachel Webster and the referee Andrea Ferrara for very useful comments on the manuscript. AL acknowledges generous support from the Institute for Advanced Study at Princeton and the John Simon Guggenheim Memorial Fellowship. This work was also supported in part by NSF grants AST-0071019, AST-0204514 and NASA grant ATP02-0004-0093.

REFERENCES

- Abel, T., Bryan, G. L., & Norman, M. L. 2002, *Science*, 295, 93
 Barkana, R. & Loeb, A. 1999, *ApJ*, 523, 54
 Barkana, R. & Loeb, A. 2001, *Physics Reports*, 349, 125
 Bell, E. F. & de Jong, R. S. 2001, *ApJ*, 550, 212
 Bromm, V., Coppi, P. S., & Larson, R. B. 2002, *ApJ*, 564, 23
 Bromm, V., Kudritzki, R. P., & Loeb, A. 2001, *ApJ*, 552, 464
 Bromm, V. & Loeb, A. 2002, *ApJ*, 575, 111
 Cen, R., 2002, *astro-ph/0210473*
 Cen, R., 2003, *astro-ph/0303236*
 Ciardi, B. & Loeb, A. 2000, *ApJ*, 540, 687
 Ciardi, B., Ferrara, A., White, S.D.M. 2003, *astro-ph/0302451*
 Dekel, A. & Silk, J. 1986, *ApJ*, 303, 39
 Dekel, A., & Woo, J. 2002, *MNRAS*, submitted; *astro-ph/0210454*
 Furlanetto, S. R. & Loeb, A. 2002, *ApJ*, 579, 1
 Furlanetto, S. R. & Loeb, A. 2003, *astro-ph/0211496*
 Haiman, Z., Rees, M. J., & Loeb, A. 1996, *ApJ*, 467, 522
 Haiman, Z., Rees, M. J., & Loeb, A. 1997, *ApJ*, 476, 458; 484, 985
 Haiman, Z., Thoul, A. A., & Loeb, A. 1996, *ApJ*, 464, 523
 Haiman, Z., Abel, T., & Rees, M. J. 2000, *ApJ*, 534, 11
 Haiman, Z., Holder, G., 2003, *astro-ph/0302403*
 Heger, A. & Woosley, S. E. 2002, *ApJ*, 567, 532
 Hui, L., Haiman, Z., 2003, *astro-ph/0302439*
 Kauffmann, G. et al., 2002, *MNRAS* accepted, *astro-ph/0205070*
 Kaplinghat, M., Chu, M., Haiman, Z., Holder, G. P., Knox, L., & Skordis, C. 2003, *ApJ*, 583, 24
 Kitayama, T. & Ikeuchi, S. 2000, *ApJ*, 529, 615
 Kogut, A. et al. 2003, *ApJ*, submitted; *astro-ph/0302213*
 Lamb, D. Q. & Reichart, D. E. 2000, *ApJ*, 536, 1
 Loeb, A. 2001, in “Supernovae and Gamma-Ray Bursters”, edited by K. W. Weiler, Springer-Verlag, in press; *astro-ph/0106455*
 Mackey, J., Bromm, V., Hernquist, L., 2002, *astro-ph/0208447*
 Miralda-Escudé, J., Haehnelt, M., & Rees, M. J. 2000, *ApJ*, 530, 1
 Nishi, R. & Tashiro, M. 2000, *ApJ*, 537, 50
 Oh, S. P., Haiman, Z. 2002, *ApJ*, 569, 558, L1
 Oh, S. P., Nollett, K. M., Madau, P., & Wasserburg, G. J. 2001, *ApJ*, 562, L1
 Ricotti, M., Gnedin, N. Y., & Shull, J. M. 2002, *ApJ*, 575, 49
 Salvaterra, R., Ferrara, A., 2003, *MNRAS*, 339, 973
 Schneider, R., Ferrara, A., Natarajan, P., Omukai, K. 2002, *ApJ*, 571, 30
 Sheth, R. K. & Tormen, G. 1999, *MNRAS*, 308, 119
 Sokasian, A., Abel, T., Hernquist, L., Springel, V., 2003, *astro-ph/0303098*
 Somerville, R.S., Livio, M., 2003, *astro-ph/0303017*
 Spergel, D. et al. 2003, *ApJ*, submitted; *astro-ph/0302209*
 Steidel, C. C., Pettini, M., & Adelberger, K. L. 2001, *ApJ*, 546, 665
 Theuns, T., Schaye, J., Zaroubi, S., Kim, T., Tzanavaris, P., & Carswell, B. 2002, *ApJ*, 567, L103
 Thoul, A.A. & Weinberg, D.H., 1996, *ApJ*, 465, 608
 Tumlinson, J. & Shull, J. M. 2000, *ApJ*, 528, L65
 Venkatesan, A., Tumlinson, J. & Shull, J.M. 2003, *ApJ*, 584, 621
 Wyithe, J. S. B. & Loeb, A. 2002a, *ApJ*, 581, 886
 Wyithe, J. S. B., & Loeb, A. 2003, *ApJ*, in press; *astro-ph/0209056*
 Zaldarriaga, M. 1997, *Phys. Rev. D*, 55, 1822